arXiv:1402.5206v1 [math.NT] 21 Feb 2014

SOLUTION OF CERTAIN PELL EQUATIONS

ZAHID RAZA, HAFSA MASOOD MALIK

ABSTRACT. Let a, b, c be any positive integers such that $c \mid ab$ and $d_i^{\pm}$ is a square free positive integer of the form $d_i^{\pm} = a^{2k}b^{2l} \pm ic^m$ where $k, l \geq m$ and $i = 1, 2$. The main focus of this paper to find the fundamental solution of the equation $x^2 - d_i^{\pm}y^2 = 1$, with the help of the continued fraction of $\sqrt{d_i^{\pm}}$. We also obtain all the positive solutions of the equations $x^2 - d_i^{\pm}y^2 = \pm 1$ and $x^2 - d_i^{\pm}y^2 = \pm 4$ by means of the Fibonacci and Lucas sequences.

Furthermore, in this work, we derive some algebraic relations on the Pell form $F_{d_i^{\pm}}(x, y) = x^2 - d_i^{\pm}y^2$ including cycle, proper cycle, reduction and proper automorphism of it. We also determine the integer solutions of the Pell equation $F_{\Delta_{d_i^{\pm}}}(x, y) = 1$ in terms of $d_i^{\pm}$.

We generalized all the results of the papers [2], [9], [26] and [37].

1. INTRODUCTION

Let d be a positive integer which is not a perfect square and N be any nonzero fixed integer. Then the equation $x^2 - dy^2 = N$ is known as Pell equation after the name of English mathematician, John Pell. The equations $x^2 - dy^2 = 1$ and $x^2 - dy^2 = -1$ are known as the classical Pell equations. If $a^2 - db^2 = N$, we say that (a, b) is a solution of the Pell equation $x^2 - dy^2 = N$. We use the notation (a, b) and $a + \sqrt{d}b$ interchangeably to denote the solutions of the equation $x^2 - dy^2 = N$. Also, if a and b are positive, we say that $a + b\sqrt{d}$ is a positive solution to the equation $x^2 - dy^2 = N$. Among these there is a least solution $a_1 + b_1\sqrt{d}$, in which a_1 and b_1 have their least positive values. Then the number $a_1 + b_1\sqrt{d}$ is called fundamental solution of the equation $x^2 - dy^2 = N$. If $a + \sqrt{d}b$ and $r + \sqrt{d}s$ are solutions of the equation $x^2 - dy^2 = N$, then $a = r$ iff $b = s$, and $a + \sqrt{d}b < r + \sqrt{d}s$ iff $a < r$ and $b < s$.

An equation $x^2 - dy^2 = 1$ has infinite many solutions iff the equation $x^2 - dy^2 = -1$ has no solution. The continued fraction of $\sqrt{d}$ played a vital role to solve the Pell equation $x^2 - dy^2 = \pm 1$. Actually its period length is useful for knowing the solution of this equation. Let d be a positive integer that is not a perfect square. Then there

1991 *Mathematics Subject Classification.* 11D09, 11D79, 11D45, 11A55, 11B39, 11B50.

Key words and phrases. The Pell equations, continued fraction, integer solutions, the generalized Fibonacci and Lucas sequences, binary quadratic form, cycle, proper cycle.

is a continued fraction expansion of $\sqrt{d}$ such that $\sqrt{d} = [a_0, \overline{a_1, a_2, \ldots, a_{n-1}, 2a_0}]$ where n is the period length and the $a_j's$ are given by the recursion formula; $a_0 = \sqrt{d}, a_k = \llcorner a_k \lrcorner$ and $a_{k+1} = \frac{1}{x_k - a_k}, k = 0, 1, 2, \ldots$

Recall that $a_n = 2a_0$ and $a_{n+k} = a_k$ for all $k \geq 1$. Then n^{th} convergent of$\sqrt{d}$ is given by

$$\frac{p_n}{q_n} = [a_0, \overline{a_1, a_2, \ldots, a_{n-1}, a_n}] = a_0 + \cfrac{1}{a_1 \cfrac{1}{a_2 + \cfrac{1}{\ddots \cfrac{1}{a_{n-1} + \cfrac{1}{a_n}}}}} \qquad n \geq 0$$

In this paper, we give the fundamental solution of the equation $x^2 - d_i^{\pm} y^2 = \pm 1$ by means of the period length of the continued fraction expansion of $\sqrt{d_i^{\pm}}$, where $d_i^{\pm} = a^{2k} b^{2l} \pm i c^m$. After finding the fundamental solution of the Pell equation $x^2 - d_i^{\pm} y^2 = \pm 1$, we obtain the positive integer solutions of equation $x^2 - d_i^{\pm} y^2 = \pm 4$, for $d_i^{\pm} = a^{2k} b^{2l} \pm i c^m$ by the means of the generalized Fibonacci and Lucas sequences. The main results of this paper also generalized the results presented in [2], [9] and [26].

Furthermore, in this work, we derive some algebraic relations on the Pell form $F_{d_i^{\pm}}(x, y) = x^2 - d_i^{\pm} y^2$ or of discriminant $\Delta_{d_i^{\pm}} = 4d_i^{\pm}$ including cycle, proper cycle, reduction and proper automorphism of it. Also we determine the integer solutions of the Pell equation $F_{\Delta_{d_i^{\pm}}}(x, y) = 1$ via $d_i^{\pm}$. The main results of this paper also generalized the results presented in [37].

2. Basic Setup

If N is a quadratic non-residue modulo d, then the Pell Equation $x^2 - dy^2 = N$ has no integer solution. If N is a perfect square, then the Pell Equation $x^2 - dy^2 = N$ is solvable in integers for all positive, non-square integers d. The equation $x^2 - dy^2 = 1$ has a solution in positive integers x and y for all positive, non-square integers d. If $(x, y) = (u, v)$ is a positive integer solution of $x^2 - dy^2 = 1$, then there exists a positive integer m such that $u + v\sqrt{d} = x_1 + y_1 m\sqrt{d}$, where (x_1, y_1) is the fundamental solution of $x^2 - dy^2 = 1$.

If we know the fundamental solutions of the equations $x^2 - dy^2 = \pm 1$ and $x^2 - dy^2 = \pm 4$ then we can give all positive integer solutions to these equations. For more information about the Pell equation, one can consult [25] and [20].

The generalized Fibonacci and Lucas sequences $f_n(w, z)$ and $L_n(w, z)$ are given in the followings:

Let w and z be two nonzero positive integers with $w^2 + 4z \geq 0$. The generalized Fibonacci and Lucas sequences with initial conditions $f_0(w, z) = 0, f_1(w, z) = 1$ and $L_0(w, z) = 2, L_1(w, z) = 0$ are of the form $f_n(w, z) = wf_{n-1}(w, z) + zf_{n-2}(w, z)$, $L_n(w, z) = wL_{n-1}(w, z) + zL_{n-2}(w, z)\ \forall\ n \geq 2$, respectively. They can also be

represented by the closed formula $f_n(w,z) = \frac{\alpha^n - \beta^n}{\alpha - \beta}$ and $L_n(w,z) = \alpha^n + \beta^n$, where $\alpha = \frac{w+\sqrt{w^2+4z}}{2}$, $\beta = \frac{w-\sqrt{w^2+4z}}{2}$. This identity is well known as Binet's formula. It is easy to see that $\alpha + \beta = w, \alpha - \beta = \sqrt{w^2+4z}$ and $\alpha\beta = -z$. For more information about the generalized Fibonacci and Lucas sequences, one can consult [16], [30], [22] and [23].

Lemma 2.1. *Let $\frac{p_k}{q_k}$ be the convergent of the continued fraction expansion of $\sqrt{d}$, and let l be the length of the expansion.*

- *If l is even, then the fundamental solution of $x^2 - dy^2 = 1$ is given by*

$$x = p_{l-1} \quad y = q_{l-1}$$

and the equation $x^2 - dy^2 = -1$ has no solutions.
- *If l is odd, then the fundamental solution of $x^2 - dy^2 = 1$ is given by*

$$x = p_{2l-1} \quad y = q_{2l-1}$$

and $x = p_{l-1}, y = q_{l-1}$ is the fundamental solution of $x^2 - dy^2 = -1$.

Theorem 2.2. *If x_1, y_1 is the fundamental solution of $x^2 - dy^2 = 1$, then every positive solution of the equation is given by x_n, y_n, where x_n and y_n are integers determined from $x_n + y_n\sqrt{d} = (x_1 + y_1\sqrt{d})^n \qquad n = 1, 2, 3, \ldots$*

Theorem 2.3. *If x_1, y_1 is the fundamental solution of $x^2 - dy^2 = -1$, then every positive solution of the equation is given by x_n, y_n, where x_n and y_n are integers determined from $x_n + y_n\sqrt{d} = (x_1 + y_1\sqrt{d})^{2n-1} \qquad n = 1, 2, 3, \ldots$*

The following two theorems are given in [20].

Theorem 2.4. *If x_1, y_1 is the fundamental solution of $x^2 - dy^2 = 4$, then every positive solution of the equation is given by x_n, y_n, where x_n and y_n are integers determined from $x_n + y_n\sqrt{d} = \frac{(x_1+y_1\sqrt{d})^n}{2^{n-1}} \qquad n = 1, 2, 3, \ldots$*

Theorem 2.5. *If x_1, y_1 is the fundamental solution of $x^2 - dy^2 = -4$, then every positive solution of the equation is given by x_n, y_n, where x_n and y_n are integers determined from $x_n + y_n\sqrt{d} = \frac{(x_1+y_1\sqrt{d})^n}{4^{n-1}} \qquad n = 1, 2, 3, \ldots$*

The following theorems are given in [32].

Theorem 2.6. *Let $d \equiv 2\ (mod 4)$ or $d \equiv 3\ (mod 4)$. Then the equation $x^2 - dy^2 = -4$ has no solution if and only if the equation $x^2 - dy^2 = -1$ has positive solutions.*

Theorem 2.7. *Let $d \equiv 0\ (mod 4)$. If $x_1 + \frac{d}{4}y_1$ is the fundamental solution of the equation $x^2 - \frac{d}{4}y^2 = 1$, then the fundamental solution of the equation $x^2 - dy^2 = 4$ is given as $(2x_1, y_1)$.*

Theorem 2.8. *Let $d \not\equiv 0\ (mod 4)$. If $x_1 + y_1\sqrt{d}$ is the fundamental solution of the equation $x^2 - dy^2 = 1$ then the fundamental solution of the equation $x^2 - dy^2 = 4$ is $(2x_1 2y_1)$.*

3. Basic Setup 2

A real binary quadratic form (or just a form) F is a polynomial in two variables x and y of the type

$$F = F(x, y) = ax^2 + bxy + cy^2 \tag{3.1}$$

with real coefficients a, b, c. We denote F briefly by $F = (a, b, c)$. The discriminant of F is defined by the formula $b^2 - 4ac$ and is denoted by Δ. A quadratic form F of discriminant Δ is called indefinite if $\Delta > 0$, and is called integral if and only if $a, b, c \in \mathbb{Z}$. An indefinite quadratic form $F = (a, b, c)$ of discriminant Δ is said to be reduced if

$$|\sqrt{\Delta} - 2|a|| < b < \sqrt{\Delta} \tag{3.2}$$

Most properties of quadratic forms can be giving by the aid of extended modular group $\overline{\Gamma}$ (see [34]). Gauss defined the group action of $\overline{\Gamma}$ on the set of forms as follows:

$$gF(x, y) = (ar^2+brs+cs^2)x^2+(2art+bru+bts+2csu)xy+(at^2+btu+cu^2)y^2 \tag{3.3}$$

for $g = \begin{pmatrix} r & s \\ t & u \end{pmatrix} \in \overline{\Gamma}$. An element $g \in \overline{\Gamma}$ is called an automorphism of F if $gF = F$. If $\det g = 1$, then g is called a proper automorphism of F and if $\det g = -1$, then g is called an improper automorphism of F. Let $Aut(F)^+$ denote the set of proper automorphisms of F and let $Aut(F)^-$ denote the set of improper automorphisms of F (for further details on binary quadratic forms see [14], [5], [7] and [24]). Let $\rho(F)$ denotes the normalization (it means that replacing F by its normalization) of $(c, -b, a)$. To be more explicit, we set

$$\rho^{j+1}(F) = (c_j, -b_j + 2c_j r_j, c_j r_j^2 - b_j r_j + a_j), \tag{3.4}$$

where

$$r_j = \begin{cases} \operatorname{sign}(c_j)\left\lfloor \frac{b_j}{2|c_j|} \right\rfloor, & \text{for } |c_j| \geq \sqrt{\Delta}; \\ \\ \operatorname{sign}(c_j)\left\lfloor \frac{b_j+\sqrt{\Delta}}{2|c_j|} \right\rfloor, & \text{for } |c_j| < \sqrt{\Delta}. \end{cases} \tag{3.5}$$

for $j \geq 0$. The number r j's called the reducing number and the form $\rho^{j+1}(F)$ is called the reduction of F. Further if F is reduced, then so is $\rho^{j+1}(F)$. In fact, ρ is a permutation of the set of all reduced indefinite forms. Let $\tau(F) = \tau(a, b, c) = (-a, b, -c)$. Then the cycle of F is the sequence $((\tau\rho)^j(G))$ for $j \in \mathbb{Z}$, where $G = (A, B, C)$ is a reduced form with $A > 0$ which is equivalent to F. The cycle and proper cycle of F is given by the following theorem [13]:

Theorem 3.1. *Let $F = (a, b, c)$ be reduced indefinite quadratic form of discriminant Δ. Then the cycle of F is a sequence $F_0 \sim F_1 \sim F_2 \sim \ldots \sim F_{l-1}$ of length l, where $F_0 = F = (a_0, b_0, c_0)$,*

$$s_j = s(F_j) = \Big\lfloor \frac{b_j + \sqrt{\Delta}}{2|c_j|} \Big\rfloor \quad \text{and} \tag{3.6}$$

$$F_{j+1} = (a_{j+1}, b_{j+1}, c_{j+1}) = (|c_{j+}|, -b_j + 2|c_j|s_j, -(c_j s_j^2 + b_j s_j + a_j)) \tag{3.7}$$

for $1 \leq i \leq l-2$. If l is odd, then the proper cycle of F is $F_0 \sim \tau F_1 \sim F_2 \sim \tau F_3 \cdots \sim \tau F_{l-2} \sim F_{l-1} \sim \tau F_0 \sim F_1 \sim \tau F_2 \sim F_3 \cdots \sim\sim F_{l-2}\tau F_{l-1}$ of length $2l$. In this case the equivalence class of F is equal to the proper equivalence class of F, and if l is even, then the proper cycle of F is $F_0 \sim \tau F_1 \sim F_2 \sim \tau F_3 \ldots \sim F_{l-2} \sim \tau F_{l-1}$ of length l. In this case the equivalence class of F is the disjoint union of the proper equivalence class of F and the proper equivalence class of $\tau(F)$.

4. Main Results

In this section, we will give our main results about the positive solutions of the Pell equations $x^2 - dy^2 = \pm 1$, and $x^2 - dy^2 = \pm 4$. for particular values of d. More precisely, for $d = d_i^{\pm} = a^{2k}b^{2l} \pm ic^m$, all the positive solutions of the equation in terms of the generalized Fibonacci and Lucas sequences has been investigated. Throughout in this section h will be a positive integer such that $h = \frac{a^k b^l}{c^m}$, because $c \mid ab$.

Theorem 4.1. *Let $h = \frac{a^k b^l}{c^m}$ be a positive integer, then the continued fraction of*

i: d_1^- *has of the form* $[a^k b^l; \overline{1, 2h-2, 1, 2a^k b^l - 2}]$ $\quad ab \geq 2$
ii: d_1^+ *has of the form* $[a^k b^l; \overline{2h, 2a^k b^l}]$
iii: d_2^- *has of the form* $[a^k b^l - 1; \overline{1, h-2, 1, 2a^k b^l - 2}]$ $\quad ab \geq 3$
iv: d_2^+ *has the form* $[a^k b^l; \overline{h, 2a^k b^l}]$

Proof. For d_1^-, the continued fraction

$$\begin{aligned}
\sqrt{a^{2k}b^{2l}-c^m} &= a^kb^l-1+\sqrt{a^{2k}b^{2l}-c^m}-(a^kb^l-1)\\
&= a^kb^l-1+\frac{(\sqrt{a^{2k}b^{2l}-c^m}-(a^kb^l-1))(\sqrt{a^{2k}b^{2l}-c^m}+a^kb^l-1)}{\sqrt{a^{2k}b^{2l}-c^m}+a^kb^l-1}\\
&= a^kb^l-1+\frac{1}{\frac{\sqrt{a^{2k}b^{2l}-c^m}+a^kb^l-1}{2a^kb^l-c^m-1}}\\
&= a^kb^l-1+\frac{1}{1+\frac{\sqrt{a^{2k}b^{2l}-c^m}-(a^kb^l-c^m)}{2a^kb^l-c^m-1}}\\
&= a^kb^l-1+\frac{1}{1+\frac{c^m}{\sqrt{a^{2k}b^{2l}-c^m}+a^kb^l-c^m}}\\
&= a^kb^l-1+\frac{1}{1+\frac{1}{\frac{\sqrt{a^{2k}b^{2l}-c^m}+a^kb^l-c^m}{c^m}}}\\
&= a^kb^l-1+\frac{1}{1+\frac{1}{2h-2+\frac{\sqrt{a^{2k}b^{2l}-c^m}-(a^kb^l-c^m)}{2a^kb^l-c^m-1}}}\\
&= a^kb^l-1+\frac{1}{1+\frac{1}{2h-2+\frac{1}{\frac{\sqrt{a^{2k}b^{2l}-c^m}+(a^kb^l-c^m)}{2a^kb^l-c^m-1}}}}\\
&= a^kb^l-1+\frac{1}{1+\frac{1}{2h-2+\frac{1}{1+\frac{\sqrt{a^{2k}b^{2l}-c^m}-(a^kb^l-1)}{2a^kb^l-c^m-1}}}}\\
&= a^kb^l-1+\frac{1}{1+\frac{1}{2h-2+\frac{1}{1+\frac{1}{\sqrt{a^{2k}b^{2l}-c^m}+(a^kb^l-1)}}}}\\
&= a^kb^l-1+\frac{1}{1+\frac{1}{2h-2+\frac{1}{1+\frac{1}{2a^kb^l-2+\sqrt{a^{2k}b^{2l}-c^m}-(a^kb^l-1)}}}}
\end{aligned}$$

Hence $\sqrt{a^{2k}b^{2l}-c^m}$ has the continued fraction of the form $[a^kb^l-1;\overline{1,2h-2,1,2a^kb^l-2}]$. Similarly for d_2^-, one can obtained the required form of the continued fraction.

For d_2^+, the continued fraction

$$\begin{aligned}\sqrt{a^{2k}b^{2l}+2c^m} &= a^kb^l+(\sqrt{a^{2k}b^{2l}+2c^m}-a^kb^l)\\ &= a^kb^l+\frac{(\sqrt{a^{2k}b^{2l}+2c^m}-a^kb^l)(\sqrt{a^{2k}b^{2l}+2c^m}+a^kb^l)}{\sqrt{a^{2k}b^{2l}+2c^m}+a^kb^l}\\ &= a^kb^l+\frac{1}{\frac{\sqrt{a^{2k}b^{2l}+2c^m}+a^kb^l}{2c^m}} = a^kb^l+\frac{1}{h+\frac{\sqrt{a^{2k}b^{2l}+2c^m}-a^kb^l}{2c^m}}\\ &= a^kb^l+\frac{1}{h+\frac{1}{\sqrt{a^{2k}b^{2l}+2c^m}-a^kb^l}} = a^kb^l\frac{1}{h+\frac{1}{2a^kb^l+\sqrt{a^{2k}b^{2l}+2c^m}-a^kb^l}}\end{aligned}$$

Hence $\sqrt{a^{2k}b^{2l}+2c^m}$ has the continued fraction of the form $[a^kb^l;\overline{h,2a^kb^l}]$. Similarly for d_1^+, one can obtained the required form of the continued fraction. □

Corollary 4.2. *If $c=1$, then the continued fraction of*

i: *d_1^- has of the form $[a^kb^l;\overline{1,2a^kb^l-2}]$*
ii: *d_1^+ has the form $[a^kb^l;\overline{2a^kb^l}]$*
iii: *d_2^- has of the form $[a^kb^l-1;\overline{1,a^kb^l-2,1,2a^kb^l-2}]$*
iv: *d_1^+ has the form $[a^kb^l;\overline{a^kb^l,2a^kb^l}]$*

Remark **4.3.** The continued fraction of $d_3=\sqrt{a^{2k}-a^k}$ is of the form $[a^k-1;\overline{2,2a^k-2}]$

Theorem 4.4. **i:** *Let us consider the Pell equation $x^2-d_1^{\pm}y^2=1$, then the fundamental solution $(x_1^{\pm1},y_1^{\pm1})$ is of the form $(2ha^kb^l\pm1,2h)$ and the other solutions are $(x_n^{\pm1},y_n^{\pm1})$, where*

$$\frac{x_n^{+1}}{y_n^{+1}}=[a^kb^l;\underbrace{2h,2a^kb^l}_{(n-1)time},2h] \qquad \textit{and}$$

$$\frac{x_n^{-1}}{y_n^{-1}}=[a^kb^l-1;\underbrace{1,2h-2,1,2a^kb^l-2}_{(n-1)time},1] \qquad \textit{respectively.}$$

ii: *Let us consider the Pell equation $x^2-d_2^{\pm}y^2=1$, the fundamental solution $(x_1^{\pm2},y_1^{\pm2})$ is of the form $(ha^kb^l\pm1,h)$ and the other solutions are $(x_n^{\pm2},y_n^{\pm2})$, where*

$$\frac{x_n^{+2}}{y_n^{+2}}=[a^kb^l;\underbrace{h,2a^kb^l}_{(n-1)time},h] \qquad \textit{and}$$

$$\frac{x_n^{-2}}{y_n^{-2}}=[a^kb^l-1;\underbrace{1,h-2,1,2a^kb^l-2}_{(n-1)time},1] \qquad \textit{respectively.}$$

Proof. The period length of the continued fraction of $\sqrt{d_i^+}$ is 2 by theorem 4.1. Since $p_{-2}=0, p_{-1}=1, q_{-2}=1, q_{-1}=0, \quad p_k=a_kp_{k-1}+p_{k-2}, q_k=a_kq_{k-1}+q_{k-2}$, therefore the fundamental solution is of the form $p_1^{+i}=a_1p_0+p_{-1}=2ha^kb^l+1, q_1^{+i}=a_1q_0+q_{-1}=2h$ for d_1^+ by using lemma 2.1. Similarly the equation $x^2-d_2^+y^2=1$ has the required solution form due to the theorem 4.1 and lemma 2.1.

Now we assume that (x_{n-1},y_{n-1}) is a solution, that is, $x_{n-1}^2-d_2^+y_{n-1}^2=1$. Then we have that

$$\frac{x_n^{+2}}{y_n^{+2}}=a^kb^l+\cfrac{1}{h+\cfrac{1}{2a^kb^l+\cfrac{1}{\ddots\ 2a^kb^l+\frac{1}{h}}}}$$

$$=a^kb^l+\cfrac{1}{h+\cfrac{1}{a^kb^l+a^kb^l+\cfrac{1}{h+\cfrac{1}{\ddots\quad 2a^kb^l+\frac{1}{h}}}}}=a^kb^l+\cfrac{1}{h+\cfrac{1}{a^kb^l+\frac{x_{n-1}}{y_{n-1}}}}=\frac{x_n}{y_n}=\frac{(ha^kb^l+1)x_{n-1}+hdy_{n-1}}{hx_{n-1}+(ha^kb^l+1)y_{n-1}}$$

$$\frac{x_n^{+2}}{y_n^{+2}}=\frac{x_1^{+2}x_{n-1}+y_1^{+2}dy_{n-1}}{y_1^{+2}x_{n-1}+x_1^{+2}y_{n-1}} \tag{4.1}$$

Similarly for d_1^+ and d_i^- we can get all positive solutions of the required form. □

Corollary 4.5. *The equation $x^2-d_i^{\pm}y^2=-1$ has no positive integer solutions, except $x^2-d_1^+y^2=-1$ has solution $(a^kb^l,1)$ only if $c=1$.*

Proof. The continued fraction of $\sqrt{d_i^{\pm}}$ have even length, therefore the equation $x^2-d_i^{\pm}y^2=-1$ has no solution by lemma 2.1, but if $c=1$, then $\sqrt{d_1^+}$ have odd length, therefore the equation has solution by lemma 2.1 and get required solution. □

Theorem 4.6. *The n^{th} integer solution $(x_n^{\pm i},y_n^{\pm i})$ of $x^2-d_i^{\pm}y^2=-1$ can be given as a linear combination of $x_i^{\pm i},y_i^{\pm i}$ and $d_i^{\pm}$ namely, for $n\geq 2$*

$$\begin{aligned} x_n^{\pm i} &= x_1^{\pm i}x_{n-1}+y_1^{\pm i}d_i^{\pm}y_{n-1} \\ y_n^{\pm i} &= y_1^{\pm i}x_{n-1}+x_1^{\pm i}y_{n-1} \end{aligned}$$

and also satisfy the recurrence relation for $n\geq 4$

$$\begin{aligned} x_n^{\pm i} &= (2x_1^{\pm i}-1)(x_{n-1}+x_{n-2})-x_{n-3} \\ y_n^{\pm i} &= (2x_1^{\pm i}-1)(y_{n-1}+y_{n-2})-y_{n-3} \end{aligned}$$

Proof. The first assertion is easily seen from 4.1. The second assertion can be proved by induction on n. □

Theorem 4.7. *All positive integer solutions of the equation $x^2-d_i^{\pm}y^2=1$ are given by*

$$(x_n^{\pm i},y_n^{\pm i})=\big(\frac{1}{2}L_n(x_1^{\pm i},-1),y_1^{\pm i}f_n(x_1^{\pm i},-1)\big) \qquad n=1,2,3,\ldots$$

Proof. Consider the Pell equation $x^2 - d_i^{\pm}y^2 = 1$, then by theorem 4.4 and theorem 2.2 all positive solution of the equation are given by $x_n^{\pm i}+y_n^{\pm i}\sqrt{d_i^{\pm}} = (x_1^{\pm i}+y_1^{\pm i}\sqrt{d_i^{\pm}})^n$

Let $\alpha_i^{\pm} = x_1^{\pm i}+y_1^{\pm i}\sqrt{d_i^{\pm}}$ and $\beta_i^{\pm} = x_1^{\pm i}-y_1^{\pm i}\sqrt{d_i^{\pm}}$. Then $\alpha_i^{\pm}+\beta_i^{\pm} = 2x_1^{\pm i}, \alpha_i^{\pm}-\beta_i^{\pm} = 2y_1^{\pm i}\sqrt{d_i^{\pm}}$ and $\alpha_i^{\pm}\beta_i^{\pm} = 1$. Therefore $x_n^{\pm i}+y_n^{\pm i}\sqrt{d_i^{\pm}} = (\alpha_i^{\pm})^n, x_n^{\pm i}-y_n^{\pm i}\sqrt{d_i^{\pm}} = (\beta_i^{\pm})^n$. Thus it follows that $x_n^{\pm i} = \frac{(\alpha_i^{\pm})^n+(\beta_i^{\pm})^n}{2} = \frac{1}{2}L_n(x_1^{\pm i}, -1)$ and $y_n^{\pm i} = \frac{(\alpha_i^{\pm})^n-(\beta_i^{\pm})^n}{2\sqrt{d_i^{\pm}}} = \frac{(\alpha_i^{\pm})^n-(\beta_i^{\pm})^n}{2y_1^{\pm i}\sqrt{d_i^{\pm}}} = y_1^{\pm i}f_n(x_1^{\pm i}, -1)$.

□

Theorem 4.8. *The fundamental solution of the Pell equation $x^2 - d_i^{\pm}y^2 = 4$ is $(x_1^{\pm i}, y_1^{\pm i}) = (2x_1^{\pm i}, 2y_1^{\pm i})$.*

Proof. We know that $a^k b^l = c^m h$ so $(a^k b^l)^2 = (c^m h)^2 \Rightarrow d_i^{\pm} = c^{2m}h^2 \pm ic^m$. we will give proof for only $d_2^+ = a^{2k}b^{2l}+2c^m$. If $c = 2s$ is even, then $d_2^+ \equiv 0(\text{mod } 4)$ and $\frac{d_2^+}{4} = 2^{2m-2}s^{2m}h^2 + 2^{m-1}s^m$. Hence by theorem 4.1 and 4.4, it follows that the equation $x^2 - (2^{2m-2}s^{2m}h^2 + 2^{m-1}s^m)y^2 = 1$ has the fundamental solution $(2^m h^2 s^m + 1, 2h)$. Then, by theorem 2.7, the fundamental solution to the equation $x^2 - d_2^+y^2 = 4$ is of the form $(2^{m+1}h^2s^m + 2, 2h\sqrt{d_2^+})$. Since $c = 2s$ and $2^m hs^m = a^k b^l$, so the the equation has the fundamental solution $(2ha^k b^l + 2 + 2h\sqrt{d_2^+})$.

Assume that c is odd. Then $d_2^+ \equiv 2, 3(\text{mod } 4)$ if $a^{2k}b^{2l}$ is even or odd respectively. Thus, by theorem 4.7 and 4.4, it follows that the fundamental solution of the equation $x^2 - (2^{2m-2}s^{2m}h^2 + 2^{m-1}s^m)y^2 = 1$ is of the form $(2^{m+1}h^2s^m + 2, 2h)$. Then, by theorem 4.4, the fundamental solution to the equation $x^2 - d_2^+y^2 = 4$ is of the form $(2ha^k b^l + 2 + 2h\sqrt{d_2^+})$. Similarly, we can proof for other values of $d_i^{\pm}$. □

Theorem 4.9. *The equation $x^2 - d_i^{\pm}y^2 = -4$ has no positive integer solution, except $x^2 - d_1^+y^2 = -4$ when $c = 1$*

Proof. Let $d_1^+ = a^{2k}b^{2l} + c^m$ and assume that c is odd. Then $d_1^+ \equiv 2, 3(\text{mod } 4)$ if $a^{2k}b^{2l}$ is even or odd respectively. Thus, by theorem 2.6 and theorem 4.4 the equation $x^2 - d_1^+y^2 = -4$ has no solution.

Now suppose that c is even and the positive integer f and g are the solution of the above equation, then $f^2 - d_1^+g^2 = -4$. But d_1^+ is even and therefore f and g are even. Since $(a^k b^l)^2 = (c^m h)^2$ therefore, $f^2 - (2^{2m-2}s^{2m}h^2 + 2^{m-1}s^m)g^2 = -4$ and implies that $(\frac{f}{2})^2 - (2^{2m-2}s^{2m}h^2 + 2^{m-1}s^m)g^2 = -1$ this is impossible by theorem 4.4. similarly, for the other equations. □

Corollary 4.10. *The equation $x^2 - d_1^+y^2 = -4$ has positive integer solutions $(2a^k b^l, 2)$ if $c = 1$.*

Theorem 4.11. *All the positive solutions of the equation $x^2 - d_i^{\pm}y^2 = 4$ are given as*

$$(x^{\pm i}, y^{\pm i}) = \big(L_n(2x_1^{\pm i}, -1), y_1^{\pm i}f_n(2x_1^{\pm i}, -1)\big) \qquad n = 1, 2, 3, \ldots$$

Proof. We know by theorem 4.8 that $2x_1^{\pm i}+2y_1^{\pm i}\sqrt{d_i^{\pm}}$ is the fundamental solution of the equation $x^2-d_i^{\pm}y^2=4$. Therefore by theorem 2.4, all positive integer solution of the equation $x^2-d_i^{\pm}y^2=4$ are given by $x_n^{\pm i}+y_n^{\pm i}\sqrt{d_i^{\pm}}=\frac{1}{2^{n-1}}\big(2x_1^{\pm i}+2y_1^{\pm i}\sqrt{d_i^{\pm}}\big)^n=2(\frac{2x_1^{\pm i}+2y_1^{\pm i}\sqrt{d_i^{\pm}}}{2})^n$. Now, let us consider $\alpha_i^{\pm}=\frac{2x_1^{\pm i}+2y_1^{\pm i}\sqrt{d_i^{\pm}}}{2}$ and $\beta_i^{\pm}=\frac{2x_1^{\pm i}-2y_1^{\pm i}\sqrt{d_i^{\pm}}}{2}$. Then $\alpha_i^{\pm}+\beta_i^{\pm}=x_1^{\pm i}, \alpha_i^{\pm}-\beta_i^{\pm}=2y_1^{\pm i}\sqrt{d_i^{\pm}}$ and $\alpha_i^{\pm}\beta_i^{\pm}=1$. Thus it is easily seen that $x_n^{\pm i}+y_n^{\pm i}\sqrt{d_i^{\pm}}=2(\alpha_i^{\pm})^n$ and $x_n^{\pm i}-y_n^{\pm i}\sqrt{d_i^{\pm}}=2(\beta_i^{\pm})^n$. Therefore $x_n^{\pm i}=(\alpha_i^{\pm})^n+(\beta_i^{\pm})^n=L_n(2x_1^{\pm i},-1)$ and $y_n^{\pm i}=\frac{(\alpha_i^{\pm})^n-(\beta_i^{\pm})^n}{\sqrt{d_i^{\pm}}}=y_1^{\pm i}\frac{(\alpha_i^{\pm})^n-(\beta_i^{\pm})^n}{2h\sqrt{d_i^{\pm}}}=y_1^{\pm i}f_n(2x_1^{\pm i},-1)$. □

Thus we can give the following corollaries.

Corollary 4.12. *If $d_1^+=a^{2k}+a^m$, then $\sqrt{d_1^+}=[a^k,\overline{2a^{k-m},2a^k}]$ and if $d_2^+=a^{2k}+2a^m$, then $\sqrt{d_2^+}=[a^k,\overline{a^{k-m},2a^k}]$*

Corollary 4.13. *If $d=a^{2k}+2a^m$, then the fundamental solution to the equation $x^2-d_2^+y^2=1$ is $x_1+y_1\sqrt{d_2^+}=a^{2k-m}+a^{k-m}\sqrt{d}$ and the equation $x^2-d_2^+y^2=-1$ has no solutions.*

Corollary 4.14. *Let $d_1^+=a^{2k}+a^m$. Then the fundamental solution to the equation $x^2-d_1^+y^2=1$ is $x_1+y_1\sqrt{d_1^+}=2a^{2k-m}+1+2a^{k-m}\sqrt{d}$ and the equation $x^2-d_1^+y^2=-1$ has no solutions*

Remark **4.15.**
- If $c=b$ and $k=l=m=1$, then the main results of [9] become the corollaries of our main results.
- If $c=a$ and $k=l=m=1$ and $b=1$, then the main results of [26] become the corollaries of our main results.

5. Main Results 2

Let us consider the matrix $M_{\pm i}$ associated with $d_i^{\pm}$ and corresponding fundamental solution $x_i^{\pm}$, $y_i^{\pm}$ as

$$M_{\pm i}=\begin{pmatrix} x_1^{\pm i} & y_1^{\pm i}d_i^{\pm} \\ y_1^{\pm i} & x_1^{\pm i}\end{pmatrix} \tag{5.1}$$

In the following theorem, we able to determine the n^{th} power of $M_{\pm i}$ which we use it later. (Here, we note that $\binom{n}{2j}=\binom{n-2}{2j}+\binom{n-2}{2j-2}+2\binom{n-2}{2j-1}$ for $j=1,2,\ldots,\frac{n-2}{2}$).

Theorem 5.1. *If $n\geq 0$, then the n^{th} power of $M_{\pm i}$ is given by*

$$M_{\pm i}^n=\begin{pmatrix} M_{11}^n & M_{12}^n \\ M_{21}^n & M_{22}^n\end{pmatrix} \qquad \textit{where}$$

a: *If n is even*

$$M_{11}^n = \sum_{j=0}^{\frac{n}{2}} \binom{n}{2j} (x_1^{\pm i})^{n-2j} (y_1^{\pm i})^{2j} (d_i^{\pm})^j = M_{22}^n$$

$$M_{12}^n = \sum_{j=0}^{\frac{n-2}{2}} \binom{n}{2j+1} (x_1^{\pm i})^{n-1-2j} (y_1^{\pm i})^{2j+1} (d_i^{\pm})^{j+1}$$

$$M_{21}^n = \sum_{j=0}^{\frac{n-2}{2}} \binom{n}{2j+1} (x_1^{\pm i})^{n-1-2j} (y_1^{\pm i})^{2j+1} (d_i^{\pm})^{j}$$

b: *If n is odd*

$$M_{11}^n = \sum_{j=0}^{\frac{n-1}{2}} \binom{n}{2j} (x_1^{\pm i})^{n-2j} (y_1^{\pm i})^{2j} (d_i^{\pm})^j = M_{22}^n$$

$$M_{12}^n = \sum_{j=0}^{\frac{n-1}{2}} \binom{n}{2j+1} (x_1^{\pm i})^{n-1-2j} (y_1^{\pm i})^{2i+1} (d_i^{\pm})^{j+1}$$

$$M_{21}^n = \sum_{j=0}^{\frac{n-}{2}} \binom{n}{2j+1} (x_1^{\pm i})^{n-1-2j} (y_1^{\pm i})^{2i+1} (d_i^{\pm})^{j}$$

Proof. (a): Here we will give the proof for d_2^+ by mathematical induction on n.
If $n = 2$, then

$M_{\pm i}^2 = \begin{pmatrix} h^2a^{2k}b^{2l} + 1 + 2ha^kb^l + h^2d_2^+ & 2hd_2^+(ha^kb^l + 1) \\ 2hd_2^+(ha^kb^l + 1) & h^2a^{2k}b^{2l} + 1 + 2ha^kb^l + h^2d_2^+ \end{pmatrix}$, *where*

$M_{11}^2 = \sum_{j=0}^{1} \binom{2}{2j} (ha^kb^l + 1)^{2-2j} h^{2j} d_2^{+j} = (ha^kb^l + 1) + h^2d_2^+ = M_{22}^2$

$M_{12}^2 = \sum_{j=0}^{0} \binom{2}{2j+1} (ha^kb^l + 1)^{1-2j} h^{2j+1} d_2^{+j+1} = 2hd(ha^kb^l + 1)$

$M_{21}^2 = \sum_{j=0}^{0} \binom{2}{2j+1} (ha^kb^l + 1)^{1-2j} h^{2j+1} d_2^{+j} = 2h(ha^kb^l + 1)$

So it is true for $n = 2$. Let us assume that it is true for $n - 2$, that is, $M_{\pm i}^{n-2} = \begin{pmatrix} M_{11}^{n-2} & M_{12}^{n-2} \\ M_{21}^{n-2} & M_{22}^{n-2} \end{pmatrix}$, where $M_{11}^{n-2} = \sum_{j=0}^{\frac{n-2}{2}} \binom{n-2}{2j} (ha^kb^l + 1)^{n-2-2j} h^{2j} (d_2^+)^j = M_{22}^{n-2}$

$M_{12}^{n-2} = \sum_{j=0}^{\frac{n-4}{2}} \binom{n-2}{2j+1} (ha^kb^l+1)^{n-3-2j} h^{2j+1} (d_2^+)^{j+1}, M_{21}^{n-2} = \sum_{j=0}^{\frac{n-4}{2}} \binom{n-2}{2j+1} (ha^kb^l+1)^{n-3-2j} h^{2j+1} (d_2^+)^j$

We will prove it for n, since $M_{\pm i}^n = M_{\pm i}^{n-2} M_{\pm i}^2$, we get

$M_{11}^{n-2}[(ha^kb^l+1)+h^2(d_2^+)]+M_{12}^{n-2}[2h(ha^kb^l+1)]=[(ha^kb^l+1)^{n-2}+\binom{n-2}{2}(ha^kb^l+1)^{n-4}h^2(d_2^+)+\cdots+\binom{n-2}{n-4}(ha^kb^l+1)^2h^{n-4}(d_2^+)^{\frac{n-4}{2}}+\cdots+h^{n-2}(d_2^+)^{\frac{n-2}{2}}\cdots][(ha^kb^l+1)+h^2(d_2^+)]+[\binom{n-2}{1}(ha^kb^l+1)^{n-3}h(d_2^+)+\cdots+\binom{n-2}{3}(ha^kb^l+1)^{n-5}h^3(d_2^+)^2+\cdots+\binom{n-2}{n-5}(ha^kb^l+1)^3h^{n-5}(d_2^+)^{\frac{n-4}{2}}+\cdots+\binom{n-2}{n-3}(ha^kb^l+1)h^{n-3}(d_2^+)^{\frac{n-2}{2}}][2h(ha^kb^l+1)]$
$=[(ha^kb^l+1)^{n-2}+\binom{n-2}{2}(ha^kb^l+1)^{n-4}h^2(d_2^+)+\cdots+\binom{n-2}{n-4}(ha^kb^l+1)^2h^{n-4}(d_2^+)^{\frac{n-4}{2}}+\cdots+h^{n-2}(d_2^+)^{\frac{n-2}{2}}\cdots][(ha^kb^l+1)+h^2(d_2^+)]+[\binom{n-2}{1}(ha^kb^l+1)^{n-3}h(d_2^+)+\cdots+\binom{n-2}{3}(ha^kb^l+1)^{n-5}h^3(d_2^+)^2+\cdots+\binom{n-2}{n-5}(ha^kb^l+1)^3h^{n-5}(d_2^+)^{\frac{n-4}{2}}+\cdots+\binom{n-2}{n-3}(ha^kb^l+1)h^{n-3}(d_2^+)^{\frac{n-2}{2}}][2h(ha^kb^l+1)]$
$=(2ha^kb^l-1)^n+\binom{n-2}{2}+1+2\binom{n-2}{1}](2ha^kb^l-1)^{n-2}4h^2(d_2^+)+\binom{n-2}{4}+\binom{n-2}{2}+2\binom{n-2}{3}(ha^kb^l+1)^{n-4}h^4(d_2^+)^2+\cdots+(ha^kb^l+1)^4h^{n-4}(d_2^+)^{\frac{n-4}{2}}$
$+[1+\binom{n-2}{n-4}+2\binom{n-2}{n-3}(ha^kb^l+1)^2h^{n-2}(d_2^+)^{\frac{n-2}{2}}+h^n(d_2^+)^{\frac{n}{2}}$
$=(ha^kb^l+1)^n+\binom{n}{2}(ha^kb^l+1)^{n-2}h^2(d_2^+)+\binom{n}{4}(ha^kb^l+1)^{n-4}h^4(d_2^+)^2+\cdots+\binom{n}{n-2}(ha^kb^l+1)^2h^{n-2}(d_2^+)^{\frac{n-2}{2}}+(h)^n(d_2^+)^{\frac{n}{2}}=\sum_{j=0}^{\frac{n}{2}}\binom{n}{2j}(ha^kb^l+1)^{n-2j}h^{2j}(d_2^+)^j=M_{12}^n$

Similarly, it can be shown that for M_{12}^n, M_{21}^n, thus $M_{\pm i}^n=\begin{pmatrix} M_{11}^n & M_{12}^n \\ M_{21}^n & M_{22}^n \end{pmatrix}$ as claimed and the other cases for $d_i^{\pm}$ can be proved similarly. □

In the following theorem, we will show that the n^{th} integer solution $(x_n^{\pm i}, y_n^{\pm i})$ of $F_{\Delta_{d_i^{\pm}}}(x,y)=1$ can be deduce via $d_i^{\pm}$.

Theorem 5.2. *The n^{th} integer solution of $F\Delta_{d_i^{\pm}}(x,y)=1$ is $(x_n^{\pm i}, y_n^{\pm i})$, where*

$$x_n^{\pm i}=\begin{cases}\sum_{j=0}^{\frac{n}{2}}\binom{n}{2j}(x_1^{\pm i})^{n-2j}(y_1^{\pm i})^{2j}(d_i^{\pm})^j, & \text{if } n \text{ is even;}\\ \sum_{j=0}^{\frac{n-1}{2}}\binom{n}{2j}(x_1^{\pm i})^{n-2j}(y_1^{\pm i})^{2j}(d_i^{\pm})^j, & \text{if } n \text{ is odd.}\end{cases}$$

$$y_n^{\pm i}=\begin{cases}\sum_{j=0}^{\frac{n-2}{2}}\binom{n}{2j+1}(x_1^{\pm i})^{n-1-2j}(y_1^{\pm i})^{2j+1}(d_i^{\pm})^j, & \text{if } n \text{ is even;}\\ \sum_{j=0}^{\frac{n-2}{2}}\binom{n}{2j}(x_1^{\pm i})^{n-1-2j}(y_1^{\pm i})^{2j+1}(d_i^{\pm})^j, & \text{if } n \text{ is odd.}\end{cases}$$

Proof. From 4.6, we can write the n^{th} solution for $d_2^+=a^{2k}b^{2l}+2c^m$ $\begin{pmatrix} x_n^{+2} \\ y_n^{+2}\end{pmatrix}=\begin{pmatrix} ha^kb^l+1 & hd_1 \\ h & ha^kb^l+1\end{pmatrix}^n\begin{pmatrix} ha^kb^l \\ h\end{pmatrix}=\begin{pmatrix} M_{11}^n+M_{12}^n \\ M_{21}^n+M_{22}^n\end{pmatrix}$ after simplification, we get the required solution. □

Now we can consider the Pell form $F_{\Delta_{d_i^{\pm}}}$ and note that this form is not reduced since $|\sqrt{4d}-1|>0$. So we can give the following theorem related to reduction of $F_{\Delta_{d_i^{\pm}}}$:

Theorem 5.3. **i:** *The reduction of* $F_{\Delta_{d_1^+}}$ *is*

$$\rho^2(F_{\Delta_{d_1^+}})=(1,2hc^m,-c^m).$$

ii: *The reduction of* $F_{\Delta_{d_2^+}}$ *is*

$$\rho^2(F_{\Delta_{d_2^+}})=(1,2hc^m,-2c^m).$$

iii: *The reduction of* $F_{\Delta_{d_2^-}}$ *is*

$$\rho^2(F_{\Delta_{d_1^-}})=(1,2hc^m-2,1-(2h-1)c^m).$$

iv: *The reduction of* $F_{\Delta_{d_1^-}}$ *is*

$$\rho^2(F_{\Delta_{d_2^-}})=(1,2hc^m-2,1-2(h-1)c^m).$$

Proof. Let $F_{\Delta_{d_1^+}}=F_{\Delta_{d_{1,0}^+}}=(1,0,-d_1^+)$. Then from (3.5), we get $r_0=0$ and hence from (3.4), we have $\rho^1(F_{\Delta_{d_1^+}})=(-d_1^+,0,1)$ which is not reduced. If we apply the reduction algorithm to $\rho^2(F_{\Delta_{d_1^+}})$ again, then we find that $r_1=a^kb^l$ and so $\rho^2(F_{\Delta_{d_1^+}})=(1,2hc^m,-c^m)$ which is reduced. Similarly, we can get the reduction for others forms. □

Now we can consider the cycle and proper cycle of $\rho^2(F_{\Delta_{d_i^{\pm}}})$.

Theorem 5.4. *Let us consider the reduction* $\rho^2(F_{\Delta_{d_i^+}})$ *of* $(F_{\Delta_{d_i^+}})$. *Then*

(1) *The cycle of* $\rho^2(F_{\Delta_{d_1^+}})$ *is* $(1,2hc^m,-c^m)\sim(c^m,2hc^m,-1)$.

(2) *The cycle of* $\rho^2(F_{\Delta_{d_1^-}})$ *is* $(1,2hc^m-2,1-(2h-1)c^m)\sim(2hc^m-c^m-1,(2h-2)c^m,-c^m)\sim(c^m,2hc^m-2c^m,-2hc^m+c^m+1)\sim(2hc^m-c^m-1,2hc^m-2,-1)$.

(3) *The cycle of* $\rho^2(F_{\Delta_{d_2^+}})$ *is* $(1,2hc^m-2,1-(2h-1)c^m)\sim(2c^m,2hc^m,-1)$.

(4) *The cycle of* $\rho^2(F_{\Delta_{d_2^-}})$ *is* $(1,2hc^m-4,1-2(h-1)c^m)\sim(2hc^m-2c^m-1,2(h-2)c^m,-2c^m)\sim(2c^m,2hc^m-4c^m,-2hc^m+2c^m+1)\sim(2hc^m-2c^m-1,2hc^m-4,-1)$.

Proof. Let $\rho^2(F_{\Delta_{d_1^+}})=\rho^2(F_{\Delta_{d_{1,0}^+}})=(1,2a^kb^l,-a^kb^l)$. Then from (3.6), we get $s_0=2h$ and thus $\rho^2(F_{\Delta_{d_{1,1}^+}})=(a^kb^l,2a^kb^l,-1)$. Again from from (3.6), we get $s_1=2a^kb^l$ and hence $\rho^2(F_{\Delta_{d_{1,2}^+}})=(1,2a^kb^l,-a^kb^l)=\rho^2(F_{\Delta_{d_1^+,0}})$. So the cycle of $\rho^2(F_{\Delta_{d_1^+}})$ is

$\rho^2(F_{\Delta_{d^+_{1,1}}}) \sim \rho^2(F_{\Delta_{d^+_{1,1}}})$. Note that $l = 2$, therefore from theorem 2.2 the proper cycle of $\rho^2(F_{\Delta_{d_1^+}})$ is $(1, 2a^kb^l, -a^kb^l)) \sim (-2a^kb^l, 2a^kb^l, 1)$ of length 2. Similarly for $\rho^2(F_{\Delta_{d_2^+}})$ and$\rho^2(F_{\Delta_{d_i^-}})$, we can obtained the required cycle. □

Corollary 5.5. *The proper cycle of* $\rho^2(F_{\Delta_{d_1^+}})$ *is* $(1, 2hc^m, -c^m) \sim (-c^m, 2hc^m, 1)$ *of length* 2.
The proper cycle of $\rho^2(F_{\Delta_{d_2^+}})$ *is* $(1, 2hc^m, -2c^m) \sim (2c^m, 2hc^m, -1)$. *of length* 2.
The proper cycle of
$\rho^2(F_{\Delta_{d_1^-}})$ *is*$\rho^2(F_{\Delta_{d_1^-}})$ *is* $(1, 2hc^m - 2, 1 - (2h - 1)c^m) \sim (-2hc^m + c^m + 1, (2h - 1)c^m, c^m) \sim (-c^m, 2hc^m - 2c^m, 2hc^m - c^m - 1) \sim (-2hc^m + c^m + 1, 2hc^m - 2, 1)$. *of length 4.*
The proper cycle of
$\rho^2(F_{\Delta_{d_2^-}})$ *is* $(1, 2hc^m - 2, 1 - 2(h - 1)c^m) \sim (-2hc^m + 2c^m + 1, 2(h - 2)c^m, 2c^m) \sim (-2c^m, 2hc^m - 4c^m, 2hc^m - 2c^m - 1) \sim (-2hc^m + 2c^m + 1, 2hc^m - 4, 1)$.*of length 4.*

Now we consider the proper automorphisms of $F_{\Delta_{d_i^\pm}}$. To get this we first consider the following representations of the action of the group Γ

$$gF_{\Delta_{d_i^\pm}} = \begin{pmatrix} x_1^{\pm i} & y_1^{\pm i} \\ y_1^{\pm i} d_i^\pm & x_1^{\pm i} \end{pmatrix}$$

Then we can give the following theorem which can be proved as in the same way that theorem 5.1 was proved.

Theorem 5.6. *Let* $d_i^\pm$ *denote non-zero square free positive integer . Then*

i: *The set of proper automorphisms of* $F_{\Delta_{d_i^\pm}}$ *is*

$$Aut^+(F_{\Delta_{d_i^\pm}}) = \{ \pm g^n_{F_{\Delta_{d_i^\pm}}} : n \in Z\}.$$

ii: *The integer solutions of* $F_{\Delta_{d_i^\pm}}(x, y) = 1$ *are* $(x_n^{\pm i}, y_n^{\pm i})$, *where*

$$\begin{pmatrix} x_n^{\pm i} \\ y_n^{\pm i} \end{pmatrix} = (g^n_{F_{\Delta_{d_i^\pm}}}) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ for } n \geq 1.$$

Remark **5.7.** If $c = a$ and $k = l = m = 1$ and $b = 1$, then the main results of [37] become the corollaries of our main results.

References

[1] Adler, A., Coury, J. E., The Theory of Numbers: A Text and Source Book of Problems, Jones and Bartlett Publishers, Boston, MA, 1995.

[2] Bilge Peker And Hasan Eenay, The Generalized Fibonacci and Lucas of The Pell Equations $x^2 - (a^2b^2 - b)y^{=}N$ and $x^2 - (a^2b^2 - 2b)y^{=}N$ arXiv:1303.1838v1.(in 2013)

[3] E. J. Barbeau. Pells Equation, New York, Springer-Verlag Inc, 2003.

[4] T. Breiteig. When is the product of two oblong numbers another oblong? Math. Mag.73(2000), 120129.
[5] D. A. Buell. Binary Quadratic Forms, Clasical Theory and Modern Computations. New York, Springer-Verlag, 1989.
[6] A. Chandoul. The Pell Equation $x^2 - Dy^2 = k^2$, Adv. in Pure Maths. 1 (2011), 1622.
[7] D. E. Flath. Introduction to Number Theory, New York etc., Wiley- Interscience Publication, 1989.
[8] H. M. Edward. Fermats Last Theorem: A Genetic Introduction to Alge- braic Number Theory. Graduate Texts in Mathematics, vol. 50, Springer- Verlag, 1977.
[9] Guney, M., Solutions of the Pell equations $x^2 - (a^2b^2 + 2b)y^2 = N$ when $N \in \{1, 4\}$, Mathematica Aeterna 2(7)2012, $629 - 638$.
[10] Ismail, M. E. H., One Parameter Generalizations of the Fibonacci and Lucas Numbers, The Fibonacci Quarterly $46 - 47(2009), 167 - 180$.
[11] M. Jacobson, H. Williams. Solving the Pell Equation, CMS Books in Mathematics. New York, Springer, 2009.
[12] Jacobson, M. J., Williams, H. C., Solving the Pell Equation, Springer,2006.
[13] Jones, J. P., Representation of Solutions of Pell Equations Using Lucas Sequences, Acta Academia Pead. Agr., Sectio Mathematicae $30(2003), 75 - 86$.
[14] J. Buchmann, U. Vollmer. Binary Quadratic Forms: An Algorithmic Approach. Berlin, Heidelberg, Springer-Verlag, 2007.
[15] H. W. Lenstra. Solving the Pell Equation. Notices Amer. Math. Soc. 49 (2) (2002), 182192.
[16] Kalman, D., Mena R., The Fibonacci Numbers exposed, Mathematics Magazine $76(2003), 167 - 181$.
[17] Keskin, R., Solutions of some quadratic Diophantine equations, Computers and Mathematics with Applications,$60(2010), 2225 - 2230$.
[18] Keskin, R., Guney, M., Positive Integer Solutions of the Pell Equation $x^2 - dy^2 = N, d \in \{k^2 \pm 4, k^2 \pm 1\}$ and$N \in \{\pm 1, \pm 4\}$ (submitted).
[19] Koninck, J., Mercier, A, 1001 Problems in Classical Number Theory, American Mathematical Society, 2007.
[20] LeVeque,W.J., Topic in Number Theory, Volume I and II,Dover Publicatios,Newyork(2002).
[21] K. Matthews. The Diophantine Equation $x^2 - Dy^2 = N, D > 0$. Exposi- tion. Math. $18(2000), 323331$.
[22] McDaniel, W.L., Diophantine Representation of Lucas Sequences, The Fibonacci Quarterly $33(1995), 58 - 63$.
[23] Melham, R., Conics Which Characterize Certain Lucas Sequences, The Fibonacci Quarterly $35(1997), 248 - 251$.
[24] R. A. Mollin. Quadratics. CRS Press, Boca Raton, New York, London, Tokyo, 1996.
[25] Nagell, T., Introduction to Number Theory, Chelsea Publishing Company, New York,1981.
[26] Peker, B., Solutions of the Pell equations $x^2 - (a^2 + 2a)y^2 = N$ via generalized Fibonacci and Lucas numbers arXiv:1304.1043vl.
[27] Peker, B., Ph.D. Thesis. Selcuk University (to appear).
[28] R. A. Mollin. Fundamental Number Theory with Applications. Second Edition Chapman & Hall CRC, Boca Raton, London, New York, 2008.
[29] Redmond, D., Number Theory: An Introduction, Markel Dekker, Inc,1996.
[30] Ribenboim, P., My Numbers, My Friends, Springer-Verlag New York,Inc., 2000.
[31] Robbins, N., Beginning Number Theory. Wm.C. Brown, Oxford, London (1993).

[32] Robertson, J. P., On D so that $x^2 - Dy^2$ represents$m$ and $-m$ and not -1, Acta Mathematica Academia Paedogogocae Nyiregyhaziensis $252009, 155 - 164$.
[33] A. Tekcan. Pell Equation $x^2 - Dy^2 = 2$, II. Ir. Math. Soc. Bull. 54 (2004) 7389.A. S. Shabani. The Proof of Two Conjectures Related to Pells Equation x2 - Dy2 = 4. Int. J. Comput. Math. Sci. $2, 1(2008), 2427$.
[34] A. Tekcan, O. Bizim, M. Bayraktar. Solving the Pell Equation using the Fundamental Element of the Field $Q(\sqrt{\Delta})$. Southeast Asian Bull. Math. $30, 2(2006), 355 - 366$.
[35] A. Tekcan. The Pell Equation $x^2 - Dy^2 = 4$. Appl. Math. Sci., Ruse $1, 58(2007), 363369$.
[36] A. Tekcan. The Number of Solutions of Pell Equations $x^2 - ky^2 = N$ and $x^2 + xy - ky^2 = N$ over Fp. Ars Comb. $102(2011), 225236$.
[37] A. Tekcan. Pell Form and Pell Equation via Oblong Numbers, Serdica Math J.39 $(2013), 37 - 52$.

Department of Mathematics,, National University of Computer and Emerging Sciences, B-Block, Faisal Town, Lahore, Pakistan.

E-mail address: zahid.raza@nu.edu.pk, hafsa.masood.malik@gmail.com